



Full Research Paper
Vol. 36, No. 4, Winter 2023, p. 393-411



Recognizing and Prioritizing Smart Solutions in the Poultry Industry based on Sustainability Criteria

A. Khadiver^{1*}, F. Mojbian², Z. Torkashvand³

Received: 11-06-2022

Revised: 05-09-2022

Accepted: 14-09-2022

Available Online: 15-02-2023

How to cite this article:

Khadivar, A., Mojbian, F., & Torkashvand, Z. (2023). Recognizing and Prioritizing Smart Solutions in the Poultry Industry based on Sustainability Criteria. *Journal of Agricultural Economics & Development* 36(4): 393-411.

DOI: [10.22067/jead.2022.76267.1130](https://doi.org/10.22067/jead.2022.76267.1130)

Abstract

Livestock and poultry production and supply is one of the significant food sectors in which more production can lead to a decrease in dependence on exports and earning foreign exchange. Poultry farming is a vital industry for sustainable food supply in all countries. In this research, intelligent applications and solutions in the poultry industry are identified and prioritized using the simultaneous evaluation of criteria and alternatives (SECA) method based on criteria representing the sustainable development. Analysis showed that eighteen principal fields of intelligent solutions are identified in the poultry industry. The weights obtained for sustainable development criteria based on the SECA method are economic (0.351), social (0.3383), and environmental (0.3065) in order of value. Economic sustainability should be most important in implementing smart solutions-based projects in the poultry industry. One of the main challenges of the agricultural sector, especially the poultry industry, is traditional production utilization which leads to the overuse of land capacity. Globalization trends, climate changes, moving from a fossil fuel-based economy to an environment-based economy, competition for land, freshwater, and labor shortage have also led to more complications in supplying nutrition. Considering the potential of smart solutions in realizing sustainable development objectives, it is suggested to focus more on the environmental aspects of poultry industry projects.

Keywords: Internet of things, Poultry industry, Sustainable development, Smart solutions, SECA method

Introduction

Due to its tight relationship with the environment, the agriculture industry has the most destructive effect on the environment (Quintero and González, 2018). In order to realize higher efficiency and greater environmental compliance, we need to identify scientific and environmental-friendly methods. Various variables and parameters are

influential in agricultural development, such as water, soil, livestock inputs, organizational services, and proper management of natural resources. One of the challenges of developing countries is the limited resources and ignorance of farmers in correctly using resources (Bani Asadi and Mehrjerdi Zare, 2010). In general, the development of the agricultural sector has various environmental effects, such as the emission of greenhouse gases, the destruction of biodiversity, pollution caused by fertilizers and pesticides, soil degradation, and increased risk to human health (DeLonge, 2016). Considering the importance and role of agriculture in the development of communities and environmental concerns on the one hand and global challenges such as food security and

1- Associate Professor, Department of Management, Alzahra University, Tehran, Iran

(*- Corresponding Author Email: a.khadivar@alzahra.ac.ir)

2- Ph.D. of Industrial Management, Department of Management and Economics, Tarbiat Modares University, Tehran, Iran

3- Master of IT Management, Department of Management, Alzahra University, Tehran, Iran

population growth on the other hand, it seems necessary to implement extensive measures to realize sustainability in agriculture (Wang, 2017). By achieving targets such as reducing poverty, guaranteeing sustainable patterns of production and consumption, taking immediate action to resolve climate change and its effects, the protecting and sustainable use of oceans, seas, and natural resources (Williams *et al.*, 2018), agriculture performs a vital responsibility in achieving the Sustainable Development Goals (SDGs) approved in September 2015 by 193 countries in order to improve the social, economic and environmental conditions of the world (GeSI, 2016).

The Internet of Things (IoT) is one of the innovations of the digital age using advanced and related technologies such as mobile and wireless communication technology, Nano-technology, identification technology based on radio waves, and smart sensor technology that can connect all objects in any time and place by anything or anyone.

One of the applications of IoT is poultry farming, which can turn a manual farm into a modern semi-automatic poultry farm. In addition, the system can be installed on the android mobile applications and help control operations such as feeding, object detection, water spraying, and gas reduction in poultry farms. The proposed system can reduce the need for human labor to feed the chickens, reduce unwanted gas, and control the temperature in the farm fully automatically. Therefore, this system reduces cost, time, workforce, and environmental pollution (Azarinfar, 2015). Another achievement of the IoT is precision livestock farming (PLF) techniques, established in the last few decades.

The world population is expected to reach 10 billion people by 2050 (United Nation Website, 2023). In order to eliminate hunger and supply the necessary food for all these 10 billion people, the current capacity of agriculture should be increased by about 70%. It is impossible to achieve this objective without relying on scientific innovations. Today, smart agriculture, which refers to the

usage of technologies like Internet of Things, sensors, location systems, robots and artificial intelligence on farms, has introduced information and communication technologies as an influential factor in the efficiency and profitability of agriculture (O'Grady and O'Hare, 2017).

Information and communication technologies (ICT) have a favorable potential for improving efficiency, effectiveness, and productivity. However, these technologies are rarely used in agriculture. Small changes in production or efficiency can significantly impact the resulting profitability (O'Grady and O'Hare, 2017). Smart agriculture can construct a homogeneous path, including sanctioned techniques and technologies. Such a path is determined through market comparison and segmentation. One of the objectives of smart agriculture is to realize diversity in technologies, network the components of the agricultural sector, and ultimately move crop and livestock production systems toward sustainable agriculture (Walter *et al.*, 2017).

Poultry farming is one of the vital industries for sustainable food supply. The implementation of a smart poultry farm (SPF) includes a smart system for automatic food feeding, water sprinklers to control the temperature of the environment, and also the use of soil mixture to reduce gas in the environment. The user can remotely control the system through the android mobile applications. The operation of this smart system, in the first place, leads to the reduction of human labor activity. Also, the development of automatic chicken-feeding devices can be very useful for the growth of the poultry industry. In existing systems, chickens are fed manually by human labor. The proposed system can replace the role of the worker in the nutrition of poultry and fulfill a semi-automatic process in the poultry industry. Also, it is very important to save and adjust the high expenses of poultry houses, including the construction cost, labor cost, fuel costs of heaters, the amount of electricity consumed by lamps and fans and etc. Relying on modern science in the development of SPF provides

the possibility of saving costs (Williams *et al.*, 2018). Smart systems help poultry farmers to control their poultry farming activities. This system can facilitate poultry management and monitoring with wireless sensors and mobile solutions. Also, environmental parameters such as temperature, light, and ammonia gas are automatically controlled (Archana and Uma, 2018).

The current study regarding the purpose is considered applied research. At the same time, it is classified in the framework of descriptive research because the researcher describes smart solutions in the poultry industry based on sustainability criteria and subsequently evaluates and prioritizes the identified components and criteria in the form of a case study in the poultry industry, especially the laying hens sector. Therefore, considering the potential of the Internet of Things technology and smart solutions in creating a new path of innovative research in the field of agriculture, as well as the increasing speed of the production of scientific resources, it is necessary to identify smart solutions in the poultry industry based on sustainability criteria. To the best of our knowledge, there has been no research on smart solutions in the poultry industry based on sustainability criteria in domestic and foreign literature. As a result, based on the new approach of simultaneous evaluation of criteria and alternatives (SECA) in multi-criteria decision-making, this research has identified smart solutions in the poultry industry and then prioritized these solutions.

Background of Study

Due to the unstable production costs and global economic uncertainty, the role of PLF in sustainable food production and processing is very important. This technique uses wireless technology to collect data through the Internet of Things. One of the goals of smart agricultural systems is providing enough data to producers and ranchers to optimize the efficiency of the agricultural system and, as a result, increase the overall performance of animals or agricultural systems. The major role of PLF is related to the optimal reduction

of losses in the entire production process (Molo *et al.*, 2009). By reducing the need for manual observations and human decisions, PLF systems facilitate the automation of these processes and reduce the time and effort required to manage large numbers of livestock. PLF systems provide real-time monitoring and livestock management. Livestock management through PLF is sometimes done as a unique livestock management approach (Halachmi *et al.*, 2019). This process allows producers to manage a larger number of animals with a reliable level of care (Smith *et al.*, 2015). Individual livestock management in large poultry farms containing thousands of birds is not always possible. However, it is possible to use PLF technology to control a subset of birds and use these inputs to assess flock health (Dalimour, 2017). According to previous studies, the review and prioritization of smart solutions in the poultry industry have not been done in any research. After reviewing the research literature, smart applications and sustainability criteria have been identified in the poultry industry, presented in Table 1.

Methodology and data

According to the review of the research literature in the field of sustainable development criteria in agriculture, all the final criteria and sub-criteria identified are presented in Table 2. It should be noted that the sustainable development criteria mentioned in agriculture are all in the class of positive criteria.

Methodology Steps

In 2022, about two thousand poultry holdings were active in the laying hens' sector in Iran. The statistical population selected in this research includes faculty members of Alzahra university and poultry industry experts on poultry industry management and smart computer applications. The statistical sample of this research is selected from among the companies active in the poultry industry based on sampling methods.

Table 1- Identifying innovative smart applications in the poultry industry

| Application | Resource | Field |
|--|--|-----------------------------------|
| Environmental monitoring systems (To control environmental inputs, including temperature, air speed, ventilation rate, substrate quality, humidity and concentration of gases such as carbon dioxide and ammonia) | (Chowdhury and Morey, 2019), (Bora <i>et al.</i> , 2020), (Bustamante <i>et al.</i> , 2012), (David <i>et al.</i> , 2015), (Jackman <i>et al.</i> , 2015), (Calvet <i>et al.</i> , 2014), (Epp, 2019), (Bordin <i>et al.</i> , 2013), (McDougal, 2018) | IoT |
| Precision feeding systems (Technology to achieve accurate food conversion rate and maintain bird weight) | (Astill <i>et al.</i> , 2020), (Park <i>et al.</i> , 2022), (Zuidhof <i>et al.</i> , 2017), (Hadinia <i>et al.</i> , 2018), (Zuidhof, 2018), (Xin and Liu, 2017) | |
| Poultry welfare monitoring systems (Technology to understand the welfare of birds in terms of temperature, humidity, etc.) | (Melor <i>et al.</i> , 2020), (Astill <i>et al.</i> , 2020), (Park <i>et al.</i> , 2022), (Sassi <i>et al.</i> , 2016) | |
| Digital imaging (Achieving movement patterns of chickens to evaluate factors related to welfare) | (Corkery <i>et al.</i> , 2013), (Marchoka <i>et al.</i> , 2013), (Silvera <i>et al.</i> , 2017), (Colles <i>et al.</i> , 2016), (Sassi <i>et al.</i> , 2016), (Vanderhasselt <i>et al.</i> , 2013) | |
| Analysis of bird sounds (Evaluation of the sound of birds as an indicator of health and welfare) | (Manteuffel <i>et al.</i> , 2004), (Fontana <i>et al.</i> , 2015), (Bright, 2008), (Carroll <i>et al.</i> , 2014), (Rizwan <i>et al.</i> , 2017) | |
| Infrared thermal imaging (Control of the chicken's health status by determining the surface temperature of objects based on infrared rays) | (Nääs <i>et al.</i> , 2014), (Shinder <i>et al.</i> , 2019) | |
| Raman spectroscopy (Imaging technique to assess the gender of the chicken embryo) | (Galli <i>et al.</i> , 2016), (Neethirajan <i>et al.</i> , 2017), (Carol <i>et al.</i> , 2014), (Galli <i>et al.</i> , 2016), (Peebles, 2018) | |
| Wearable sensors for the detection of avian influenza virus (Clinical symptom detection sensor and quick virus diagnosis) | (Neethirajan <i>et al.</i> , 2017), (Okada <i>et al.</i> , 2009), (Okada <i>et al.</i> , 2014) | |
| Avian influenza virus biosensors (Biological receptor to detect the presence of a pathogen, protein, nucleic acid, etc.) | (Astill <i>et al.</i> , 2020), (Nuñez and Ross, 2019), (Luka <i>et al.</i> , 2015), (Chen and Neethirajan, 2015) | |
| Internet of things and smart poultry farming | (Park <i>et al.</i> , 2022), (Banhazi, 2009), (Bello and Zeadally, 2015), (Zuidhof <i>et al.</i> , 2017) | |
| Clustering to monitor the growth status of chickens and real-time disease diagnosis | (Aengwanich <i>et al.</i> , 2012), (Ghufran Ahmed <i>et al.</i> , 2021) | Data mining |
| PLF technology and data | (Astill <i>et al.</i> , 2020), (Banhazi, 2009) | |
| Data collection and storage | (Banhazi, 2009), (Schuetz <i>et al.</i> , 2018), (Smith <i>et al.</i> , 2015), (Chen <i>et al.</i> , 2014), (Wolfert <i>et al.</i> , 2017) | |
| Data access for smart poultry management systems | (Bumanis <i>et al.</i> , 2022), (Davis, 2016) | |
| Data governance | (Information Builders, 2011), (Wizman <i>et al.</i> , 2018), (Saykota, 2016) | |
| Big data analysis systems in the poultry industry | (Sicular, 2013), (Kamilaris <i>et al.</i> , 2017), (Wolffort <i>et al.</i> , 2017), (Chen <i>et al.</i> , 2014), (Manika <i>et al.</i> , 2011) | RFID Tags |
| Tracking the chickens in poultry halls to determine the time of rest, the time of feeding, etc. | Ronald and Sjarhei, 2012), (Rani and Devarajan, 2012), (Praveen and Satish, 2012), (Zhang <i>et al.</i> , 2007) | |
| Mobile management system and farm management system to transfer and receive farm environmental information | Chakchai <i>et al.</i> , 2014 | Mobile technology and GPS mapping |

Note: a. abbreviation: Radio-frequency identification (RFID)

Table 2- The final criteria of sustainable development

| The main criterion | Sub criterion | Resource |
|--------------------|-----------------------------|--|
| Economical | Productivity | (Veisi <i>et al.</i> , 2016; Chiou <i>et al.</i> , 2005; Quaddus and Siddique, 2001; Rezaei Moghaddam and Karami, 2008) |
| | Profitability | (Rezaei-Moghaddam and Karami, 2008; Quaddus and Siddique, 2001) |
| | Employment | (Senoret <i>et al.</i> , 2022), (Rezaei-Moghaddam and Karami, 2008) |
| Social | Quality of life | (Comim and Hirai, 2022), (Rezaei-Moghaddam and Karami, 2008) |
| | Fairness | (Quaddus and Siddique, 2001; Rezaei-Moghaddam and Karami, 2008) |
| | Partnership | (Rezaei-Moghaddam and Karami, 2008) |
| Environmental | Environmental protection | (Gunnarsdottir <i>et al.</i> , 2022), (Comim and Hirai, 2022), (Rezaei-Moghaddam and Karami, 2008); (Zarei, Mohammadian and Ghasemi, 2016) |
| | Reasonable use of resources | (Rezaei-Moghaddam and Karami, 2008) |
| | Quality of products | (Bordin <i>et al.</i> , 2022), (Rezaei-Moghaddam and Karami, 2008; Poursaeed <i>et al.</i> , 2010) |

A total of 40 questionnaires have been distributed among 20 experts. The first 20 questionnaires have been distributed to identify the components and change and remove some components. The information from the second batch of questionnaires has been used to prioritize alternatives and criteria through the SECA method. This method was presented by Keshavarz-Ghorabae *et al.* (2018) in research entitled "simultaneous evaluation of criteria and alternatives in multi-criteria decision making." The purpose of this method is to determine the total score of the alternatives and the weight of the criteria at the same time. To achieve this goal, a multi-objective nonlinear mathematical model is formulated.

Research Steps

In order to identify and prioritize smart solutions in the poultry industry based on sustainability criteria, a literature review and a study of references and background papers have been studied. As Figure 1 illustrates the steps of methodology including i) extracting criteria from literature review and interviewing with expert; ii) the relevant components and indicators were identified and finalized through consultation with experts; iii) subsequently, the importance score and weight of the criteria have been calculated with the help of poultry industry experts through the SECA method and Finally, iv) their prioritization and evaluation have been

completed (Keshavarz-Ghorabae *et al.*, 2018).

SECA Method

The steps to implement the SECA method proposed by Keshavarz-Ghorabae *et al.* (2018) are as follows:

First, the $n \times m$ decision matrix, including n alternatives and m criteria, is prepared as follows.

$$X = \begin{bmatrix} x_{11} & \dots & x_{1m} \\ \vdots & \ddots & \vdots \\ x_{n1} & \dots & x_{nm} \end{bmatrix}$$

Where X_{ij} is the evaluation of the i^{th} alternative concerning the j^{th} criterion.

Then the decision matrix is normalized based on the following relations:

$$X^N = \begin{bmatrix} x_{11}^N & \dots & x_{1m}^N \\ \vdots & \ddots & \vdots \\ x_{n1}^N & \dots & x_{nm}^N \end{bmatrix}$$

$$x_{ij}^N = \begin{cases} \frac{x_{ij}}{\max_k x_{kj}} & \text{if } j \in BC \\ \frac{\min_k x_{kj}}{x_{ij}} & \text{if } j \in NC \end{cases}$$

Where BC includes profit-focused criteria (or positive criteria), and NC includes cost-focused criteria (or negative criteria).




Figure 1- The research procedure

The standard deviation of the elements of each vector can provide the information of intra-criteria. The correlation between each pair of criteria vectors in the decision matrix is calculated to obtain information on inter-criteria. This correlation is denoted by r_{ji} . The following relationship can show the difference between the j^{th} criterion and other criteria.

$$\pi_j = \sum_{i=1}^m (1 - r_{ji}) \quad (1)$$

Increasing the variance in the vector of a criterion (j) and increasing the difference between criterion j and other criteria increases the importance (weight) of the criterion. Accordingly, the normalized values of σ_j^N and π_j^N are defined as reference points for the weights of the criteria. These values can be calculated as follows:

$$\sigma_j^N = \frac{\sigma_j}{\sum_{i=1}^m \sigma_i} \quad (2)$$

$$\pi_j^N = \frac{\pi_j}{\sum_{i=1}^m \pi_i} \quad (3)$$

In light of the above, a nonlinear multi-objective programming model is obtained as follows:

$$\max S_i = \sum_{j=1}^m w_j x_{ij}^N, \quad \forall_i \in \{1,2,3,\dots, n\} \quad (4)$$

$$\min \lambda_b = \sum_{j=1}^m (w_j - \sigma_j^N)^2 \quad (5)$$

$$\min \lambda_c = \sum_{j=1}^m (w_j - \pi_j^N)^2 \quad (6)$$

$$\text{s.t } \sum_{j=1}^m w_j = 1 \quad (7)$$

$$w_j \leq 1, \quad \forall_i \in \{1.2.3 \dots m\} \quad (8)$$

$$w_j \geq 0, \quad \forall_i \in \{1.2.3 \dots m\} \quad (9)$$

Equation 4 increases the overall performance of each alternative. Also, equations 5 and 6 minimize the weight criteria deviation from each criterion's reference points. Equation 7 guarantees that the sum of

the weights is equal to 1. Equations 8 and 9 determine the weights of the criteria for some values in the interval $[\varepsilon, 1]$. It should be said that ε is a small positive parameter considered a lower bound for the criterion weight. In this method, the value of this parameter is set equal to 0.003. To optimize the above model, we can convert the objective function into a constraint. A single-objective relationship is formulated as follows.

$$\text{Max } Z = \lambda_a - \beta(\lambda_b + \lambda_c) \tag{10}$$

$$\text{s.t } \lambda_a \leq S_i, \forall_i \in \{1,2,3,\dots,n\} \tag{11}$$

$$S_i = \sum_{j=1}^m w_j x_{ij}^N = 1, \forall_i \in \tag{12}$$

$$\{1.2.3 \dots n\}$$

$$\lambda_a = \sum_{j=1}^m (w_j - \sigma_j^N)^2 \tag{13}$$

$$\lambda_a = \sum_{j=1}^m (w_j - \Pi_j^N)^2 \tag{14}$$

$$\sum_{j=1}^m w_j = 1 \tag{15}$$

$$w_j \leq 1, \forall_i \in \{1,2,3,\dots,m\} \tag{16}$$

$$w_j \geq \varepsilon, \forall_i \in \{1,2,3,\dots,m\} \tag{17}$$

According to the objective function of the model above, the minimum overall performance score of the alternatives is maximized. Since the deviation from the reference points must be minimal, their differences from the objective function are

calculated with the coefficient B. This coefficient affects the importance of achieving reference points of weight criteria. The overall performance score of each alternative (S_i) and the weight of each criterion (w_j) are determined by solving this model. Model formulation and calculations have been done in Lingo software.

Data Acquisition

Primary data, including smart solutions and sustainable development criteria in the poultry industry, have been extracted and listed in the Table 3.

Using the questionnaire, extracting options related to smart solutions in the poultry industry will be prioritized based on sustainable development criteria in agriculture.

Results

The proposed model can simultaneously determine the overall performance score of the alternatives and the objective weight of the poultry industry's criteria. In order to verify the SECA method, the objective weight of the criteria and the overall performance of the resulting alternatives are analyzed.

Table 3- Related alternatives to smart solutions in the poultry industry

| Alternatives | Symbol |
|---|--------|
| Environmental monitoring systems | A1 |
| Precision feeding systems | A2 |
| Welfare monitoring systems | A3 |
| Digital imaging | A4 |
| Analysis of bird sounds | A5 |
| Infrared thermal imaging | A6 |
| Raman spectroscopy | A7 |
| Wearable sensors for the detection of avian influenza virus | A8 |
| Avian influenza virus biosensors | A9 |
| Internet of things and smart poultry farming | A10 |
| Clustering to monitor the growth status of chickens and real-time disease diagnosis | A11 |
| PLF technology and data | A12 |
| Data collection and storage | A13 |
| Data access for smart poultry management systems | A14 |
| Data governance | A15 |
| Big data analysis systems in the poultry industry | A16 |
| Tracking the chickens in poultry halls Using RFID tags | A17 |
| Mobile management system and GPS mapping | A18 |

Table 4- Related alternatives to smart solutions in the poultry industry

| Attributes | Symbol |
|-----------------------------|--------|
| Productivity | C1 |
| Profitability | C2 |
| Employment | C3 |
| Quality of Life | C4 |
| Fairness | C5 |
| Partnership | C6 |
| Environmental protection | C7 |
| Reasonable use of resources | C8 |
| Quality of products | C9 |

The results show that determining the appropriate value for the component (β) facilitates the determination of the sustainability weight for the criteria and performance scores of the alternatives. Finally, the results of the SECA method are compared with the results of the SD and CRITIC methods.

Related alternatives to smart solutions in the poultry industry are listed in Table 4.

In this section, final model is executed using the normalized decision matrix table data and various values for the coefficient $\beta = (0.1, 0.2, 0.3, 0.4, 0.5, 1, 2, 3, 4, 5)$. After execution of the model, ten sets of weights for the criterion are obtained. The different weight values of the criteria resulting from the change of β value are shown in Table 3. Figure 2 also shows the variation of these weights.

Table 3- Different weight values of the criteria resulting from changing the value of β

| | β | | | | | | | | | |
|-----------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 1 | 2 | 3 | 4 | 5 |
| W1 | 0.1273 | 0.1550 | 0.1571 | 0.1539 | 0.1517 | 0.1404 | 0.1339 | 0.1278 | 0.1247 | 0.1228 |
| W2 | 0.1281 | 0.1197 | 0.1208 | 0.1217 | 0.1225 | 0.1162 | 0.1097 | 0.1083 | 0.1077 | 0.1072 |
| W3 | 0.2751 | 0.2368 | 0.2001 | 0.1789 | 0.1661 | 0.1375 | 0.1271 | 0.1200 | 0.1164 | 0.1142 |
| W4 | 0.2743 | 0.2579 | 0.2158 | 0.2025 | 0.1952 | 0.1617 | 0.1311 | 0.1230 | 0.1191 | 0.1166 |
| W5 | 0.0246 | 0.0783 | 0.0939 | 0.1123 | 0.1240 | 0.1283 | 0.1215 | 0.1159 | 0.1131 | 0.1113 |
| W6 | 0.0010 | 0.0010 | 0.0010 | 0.0010 | 0.0010 | 0.0498 | 0.0880 | 0.0994 | 0.1049 | 0.1085 |
| W7 | 0.1457 | 0.1188 | 0.1281 | 0.1326 | 0.1351 | 0.1325 | 0.1284 | 0.1259 | 0.1245 | 0.1238 |
| W8 | 0.0010 | 0.0010 | 0.0362 | 0.0479 | 0.0542 | 0.0770 | 0.0963 | 0.1019 | 0.1046 | 0.1064 |
| W9 | 0.0230 | 0.0316 | 0.0469 | 0.0492 | 0.0501 | 0.0565 | 0.0640 | 0.0778 | 0.0850 | 0.0892 |

Table 4- Ranking criteria according to different values of β

| rank | β | | | | | | | | | |
|-----------|---------|-----|-----|-----|-----|---|---|---|---|---|
| | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 1 | 2 | 3 | 4 | 5 |
| W1 | 5 | 3 | 3 | 3 | 3 | 2 | 1 | 1 | 1 | 2 |
| W2 | 4 | 4 | 5 | 5 | 6 | 6 | 6 | 6 | 6 | 7 |
| W3 | 1 | 2 | 2 | 2 | 2 | 3 | 4 | 4 | 4 | 4 |
| W4 | 2 | 1 | 1 | 1 | 1 | 1 | 2 | 3 | 3 | 3 |
| W5 | 6 | 6 | 6 | 6 | 5 | 5 | 5 | 5 | 5 | 5 |
| W6 | 8 | 8 | 9 | 9 | 9 | 9 | 8 | 8 | 7 | 6 |
| W7 | 3 | 5 | 4 | 4 | 4 | 4 | 3 | 2 | 2 | 1 |
| W8 | 8 | 8 | 8 | 8 | 7 | 7 | 7 | 7 | 8 | 8 |
| W9 | 7 | 7 | 7 | 7 | 8 | 8 | 9 | 9 | 9 | 9 |




Figure 2- Variation of criteria weight according to different values of β

As shown in Figure 2, when the values of β are greater than 3 ($\beta \geq 3$), the criteria weights are more sustainable.

Now, to verify the results, the criteria weights are also determined using other methods. Two methods, SD and CRITIC, have been chosen for comparative analysis. The

weights of the criteria obtained from these three methods are shown in Table 5. If the correlation value is greater than 0.6, there is a reasonable link between the results (Walters, 2009). Table 6 shows the correlation values between the results.

Table 5- Comparing the objective weight of criteria with other methods

| | STD | CRITIC |
|----|--------|--------|
| W1 | 0.1022 | 0.1155 |
| W2 | 0.1082 | 0.1104 |
| W3 | 0.1020 | 0.1178 |
| W4 | 0.1084 | 0.1073 |
| W5 | 0.0883 | 0.1295 |
| W6 | 0.1259 | 0.1187 |
| W7 | 0.1122 | 0.1121 |
| W8 | 0.1264 | 0.1041 |
| W9 | 0.1266 | 0.0845 |

Table 6- Correlation between the weights of the criteria according to the values of β

| | STD | CRITIC |
|------------|---------|--------|
| 0.1 | -0.4168 | 0.1174 |
| 0.2 | -0.5770 | 0.2065 |
| 0.3 | -0.6275 | 0.2046 |
| 0.4 | -0.6889 | 0.2410 |
| 0.5 | -0.7273 | 0.2676 |
| 1 | -0.7805 | 0.4223 |
| 2 | -0.7594 | 0.6431 |
| 3 | 0.6903 | 0.6596 |
| 4 | 0.6145 | 0.6567 |
| 5 | 0.5272 | 0.6477 |

As shown in Table 6, the correlation values for $\beta \geq 3$ are greater than 0.6 (these values are highlighted in bold in the table). Likewise, $\beta=3$ can be a suitable threshold value for performing calculations in the proposed method.

Now, the overall performance score of each criterion is calculated by the proposed model based on the normalized decision matrix table, as well as β values and, subsequently, the weight of the obtained criteria. The calculated scores of the alternatives' overall performance

and correlation ranking are presented in Tables 7 and 8, respectively. Additionally, performance scores are visualized in Figure 3.

Table 7- Overall performance scores of alternatives with different values of β

| | β | | | | | | | | | |
|------------|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 1 | 2 | 3 | 4 | 5 |
| S1 | 0.683 | 0.671 | 0.652 | 0.647 | 0.644 | 0.627 | 0.612 | 0.603 | 0.598 | 0.596 |
| S2 | 0.653 | 0.650 | 0.628 | 0.618 | 0.612 | 0.578 | 0.553 | 0.539 | 0.532 | 0.528 |
| S3 | 0.636 | 0.629 | 0.615 | 0.610 | 0.606 | 0.582 | 0.564 | 0.553 | 0.548 | 0.545 |
| S4 | 0.709 | 0.716 | 0.690 | 0.681 | 0.677 | 0.639 | 0.609 | 0.592 | 0.584 | 0.579 |
| S5 | 0.734 | 0.753 | 0.727 | 0.724 | 0.723 | 0.681 | 0.641 | 0.624 | 0.615 | 0.609 |
| S6 | 0.657 | 0.674 | 0.656 | 0.651 | 0.649 | 0.615 | 0.586 | 0.570 | 0.563 | 0.558 |
| S7 | 0.674 | 0.678 | 0.656 | 0.649 | 0.645 | 0.614 | 0.589 | 0.576 | 0.569 | 0.565 |
| S8 | 0.796 | 0.784 | 0.752 | 0.740 | 0.733 | 0.689 | 0.656 | 0.639 | 0.631 | 0.626 |
| S9 | 0.722 | 0.722 | 0.695 | 0.686 | 0.682 | 0.642 | 0.610 | 0.594 | 0.586 | 0.581 |
| S10 | 0.652 | 0.650 | 0.631 | 0.627 | 0.625 | 0.595 | 0.569 | 0.557 | 0.551 | 0.547 |
| S11 | 0.672 | 0.680 | 0.667 | 0.661 | 0.657 | 0.630 | 0.610 | 0.598 | 0.592 | 0.588 |
| S12 | 0.750 | 0.758 | 0.727 | 0.718 | 0.714 | 0.674 | 0.640 | 0.623 | 0.615 | 0.609 |
| S13 | 0.636 | 0.639 | 0.622 | 0.617 | 0.615 | 0.583 | 0.556 | 0.543 | 0.536 | 0.532 |
| S14 | 0.642 | 0.640 | 0.616 | 0.610 | 0.606 | 0.575 | 0.550 | 0.536 | 0.530 | 0.526 |
| S15 | 0.672 | 0.673 | 0.671 | 0.671 | 0.671 | 0.650 | 0.634 | 0.625 | 0.621 | 0.618 |
| S16 | 0.636 | 0.629 | 0.615 | 0.610 | 0.606 | 0.575 | 0.550 | 0.548 | 0.548 | 0.548 |
| S17 | 0.636 | 0.629 | 0.615 | 0.610 | 0.606 | 0.575 | 0.550 | 0.536 | 0.530 | 0.526 |
| S18 | 0.636 | 0.629 | 0.615 | 0.610 | 0.606 | 0.616 | 0.626 | 0.624 | 0.622 | 0.621 |

Table 8- Ranking the overall performance scores of alternatives according to different values of β

| | β | | | | | | | | | |
|-----|---------|-----|-----|-----|-----|----|----|----|----|----|
| | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 1 | 2 | 3 | 4 | 5 |
| A1 | 6 | 10 | 10 | 10 | 10 | 8 | 6 | 6 | 6 | 6 |
| A2 | 11 | 11 | 12 | 12 | 13 | 15 | 15 | 16 | 16 | 16 |
| A3 | 14 | 15 | 17 | 17 | 14 | 14 | 13 | 13 | 14 | 14 |
| A4 | 5 | 5 | 5 | 5 | 5 | 6 | 9 | 9 | 9 | 9 |
| A5 | 3 | 3 | 3 | 2 | 2 | 2 | 2 | 3 | 4 | 4 |
| A6 | 10 | 8 | 8 | 8 | 8 | 10 | 11 | 11 | 11 | 11 |
| A7 | 7 | 7 | 9 | 9 | 9 | 11 | 10 | 10 | 10 | 10 |
| A8 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| A9 | 4 | 4 | 4 | 4 | 4 | 5 | 7 | 8 | 8 | 8 |
| A10 | 12 | 12 | 11 | 11 | 11 | 12 | 12 | 12 | 12 | 13 |
| A11 | 9 | 6 | 7 | 7 | 7 | 7 | 8 | 7 | 7 | 7 |
| A12 | 2 | 2 | 2 | 3 | 3 | 3 | 3 | 5 | 5 | 5 |
| A13 | 14 | 14 | 13 | 13 | 12 | 13 | 14 | 15 | 15 | 15 |
| A14 | 13 | 13 | 14 | 14 | 14 | 18 | 16 | 17 | 17 | 17 |
| A15 | 8 | 9 | 6 | 6 | 6 | 4 | 4 | 2 | 3 | 3 |
| A16 | 16 | 15 | 15 | 14 | 14 | 16 | 16 | 14 | 13 | 12 |
| A17 | 16 | 15 | 17 | 14 | 14 | 16 | 16 | 17 | 17 | 18 |
| A18 | 16 | 15 | 15 | 17 | 14 | 9 | 5 | 4 | 2 | 2 |

Table 9- Spearman's correlation coefficient of the resulting ranks

| | β | | | | | | | | | |
|-----|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | 0.1 | 0.2 | 0.3 | 0.4 | 0.5 | 1 | 2 | 3 | 4 | 5 |
| 0.1 | 1.000 | 0.964 | 0.947 | 0.938 | 0.938 | 0.843 | 0.759 | 0.672 | 0.619 | 0.613 |
| 0.2 | 0.964 | 1.000 | 0.977 | 0.969 | 0.963 | 0.836 | 0.722 | 0.640 | 0.596 | 0.590 |
| 0.3 | 0.947 | 0.977 | 1.000 | 0.986 | 0.986 | 0.894 | 0.782 | 0.720 | 0.679 | 0.673 |
| 0.4 | 0.938 | 0.969 | 0.986 | 1.000 | 0.990 | 0.870 | 0.735 | 0.673 | 0.622 | 0.614 |
| 0.5 | 0.938 | 0.963 | 0.986 | 0.990 | 1.000 | 0.915 | 0.793 | 0.729 | 0.679 | 0.668 |
| 1 | 0.843 | 0.836 | 0.894 | 0.870 | 0.915 | 1.000 | 0.959 | 0.926 | 0.896 | 0.885 |
| 2 | 0.759 | 0.722 | 0.782 | 0.735 | 0.793 | 0.959 | 1.000 | 0.976 | 0.959 | 0.948 |
| 3 | 0.672 | 0.640 | 0.720 | 0.673 | 0.729 | 0.926 | 0.976 | 1.000 | 0.992 | 0.987 |
| 4 | 0.619 | 0.596 | 0.679 | 0.622 | 0.679 | 0.896 | 0.959 | 0.992 | 1.000 | 0.997 |
| 5 | 0.613 | 0.590 | 0.673 | 0.614 | 0.668 | 0.885 | 0.948 | 0.987 | 0.997 | 1.000 |

As shown in Figure 3 and Table 7, when the value of β is greater than 3 ($\beta \geq 3$), the performance of alternatives is more distinct and stable.

In order to check the sustainability of the ranking criteria in different values of β , the Spearman correlation coefficient of the rankings in each column of Table 8 is calculated. The results are reflected in Table 9. As shown in Table 9, when the values of β are greater than 1, the ranks have complete sustainability. It can be said that $\beta=3$ is a suitable threshold value for determining the overall performance score and ranking of alternatives.

The results show that determining the appropriate value for the coefficient (β) facilitates realizing the sustainability weight for the criteria and performance scores for the alternatives. Table 10 shows the weight of sustainability criteria for $\beta=3$.

Table 11 shows the prioritization of alternatives based on the resulting performance scores.

Smart solutions in the poultry industry in order of priority are rapid diagnosis/point of care diagnosis, smart systems for poultry management, analysis of bird sounds, mobile technology, GPS mapping, sensors, and new technologies in poultry operations, environmental monitoring systems, communication between sensors and used equipment, wearable sensors to detect avian influenza viruses, digital imaging, Raman spectroscopy, infrared thermal imaging, biosensors for detection of avian influenza virus, welfare monitoring systems, data privacy and security, distributed data storage systems, precision feeding systems, poultry tracking using RFID tags, and other data storage systems such as cloud-based operating systems and hybrid storage systems (Table 11).




Figure 3- Variability in overall performance scores associated with β value

| Table 10- Prioritizing sustainability criteria in the poultry industry based on the SECA method for ($\beta=3$) | | | |
|---|-----------------------------|--------------------|---------------------|
| Criteria symbol | Criteria | Weight of criteria | Ranking of criteria |
| C1 | Productivity | 0.1278 | 1 |
| C 2 | Profitability | 0.1083 | 6 |
| C 3 | Employment | 0.1200 | 4 |
| C 4 | Quality of Life | 0.1230 | 3 |
| C 5 | Fairness | 0.1159 | 5 |
| C 6 | Partnership | 0.0994 | 8 |
| C 7 | Environmental protection | 0.1259 | 2 |
| C 8 | Reasonable use of resources | 0.1019 | 7 |
| C 9 | Quality of products | 0.0778 | 9 |

| Alternatives symbol | Alternatives | Weight of alternatives | Ranking |
|---------------------|--|------------------------|---------|
| A1 | Environmental monitoring systems | 0.603 | 6 |
| A2 | Precision feeding systems | 0.539 | 16 |
| A3 | Poultry welfare monitoring systems | 0.553 | 13 |
| A4 | Digital imaging | 0.592 | 9 |
| A5 | Analysis of bird sounds | 0.624 | 3 |
| A6 | Infrared thermal imaging | 0.570 | 11 |
| A7 | Raman spectroscopy | 0.576 | 10 |
| A8 | Rapid diagnosis/point of care diagnosis | 0.639 | 1 |
| A9 | Wearable sensors for the detection of avian influenza virus | 0.594 | 8 |
| A10 | Avian influenza virus biosensors | 0.557 | 12 |
| A11 | Communication between sensors and equipment used | 0.598 | 7 |
| A12 | Sensors and new technologies in poultry operations | 0.623 | 5 |
| A13 | Distributed data storage systems | 0.543 | 15 |
| A14 | Other data storage systems, such as cloud-based operating systems and hybrid storage systems | 0.536 | 17 |
| A15 | Smart systems for poultry management | 0.625 | 2 |
| A16 | Data privacy and security | 0.548 | 14 |
| A17 | Poultry tracking using RFID tags | 0.536 | 17 |
| A18 | Mobile technology and GPS mapping | 0.624 | 4 |

Conclusion

Over the past few decades, various methods for multi-criteria decision-making have been proposed. Most of these methods evaluate several alternatives based on a default set of criteria weights. In addition, there are methods to determine the objective and subjective weight of the criteria. In this study, a new approach was introduced for applying the method of simultaneous evaluation of criteria and alternatives (SECA). Subsequently, a nonlinear multi-objective mathematical model was formulated based on the introduced approach. The objective function of the model seeks to maximize the overall performance of each alternative according to the diversity of intra-criteria and inter-criteria information and the decision matrix. The results show that determining the appropriate value for the coefficient (β) can facilitate the determination

of sustainability weights for criteria and performance scores for alternatives. In the research process, smart solutions in the poultry industry were first identified based on the SECA method. Based on the analysis, 18 main areas of smart solutions in the poultry industry were determined. The identified innovative applications were prioritized based on sustainable development criteria in the next step.

The weights obtained for sustainable development criteria based on the SECA method are economic (0.351), social (0.3383), and environmental (0.3065) in order of value. Economic sustainability should be most important in implementing smart solutions-based projects in the poultry industry. One of the main challenges of the agricultural sector, especially the poultry industry, is traditional production utilization which leads to the

overuse of land capacity. Also, the globalization trends, climate changes, moving from a fossil fuel-based economy to an environment-based economy, competition for land, freshwater, and labor shortage have led to more complications in supplying nutrition.

Considering the potential of smart solutions in realizing sustainable development objectives, it is suggested to focus more on the environmental aspects of poultry industry projects.

References

1. Archana, M.P., & Uma, S.K. (2018). Monitoring and controlling of poultry farm using IOT, *International Journal of Innovation Research in Computer and Communication Engineering* 6(4).
2. Aengwanich, W., Intarakhamhaeng, M., Wandee, J., Nongbua, T., Chaiyasak, S., Srikot, P., Thammasar, K., Junsanitsri, N., Sritongtuam, K., & Tawinwaang, T. (2012). Poultry Production Clusters (PPCs) after AI Outbreaks in Thailand: Past, Present and Future Direction, *International Journal of Poultry Science* 11(8): 541-550.
3. Astill, J., Dara, R.A., Fraser, E.D.G., & Sharif, S. (2020). Smart poultry management: smart sensors, big data, and the internet of thing, *Computers and Electronics in Agriculture* 170: 105291.
4. Banhazi, T. (2009). Precision Livestock farming: a suite of electronic systems to ensure the application of best practise management on livestock farms. *Aust. Journal Multi-disciplinary Engineering* 7. <https://doi.org/10.1080/14488388.2009.11464794>.
5. Baniyasi, M., & Zare Mehrjerdi, M.R. (2010). Studying the effects of optimal cultivation pattern on rural poverty: case study of Orzoo'iyeh District in Baft (Kerman – Iran), *Journal of Agricultural Economics* 4(2): 183-209.
6. Bello, O., & Zeadally, S. (2015). Intelligent device-to-device communication in the internet of things. *IEEE Syst. Journal* 10: 1–11. <https://doi.org/10.1109/JSYST.2014.2298837>.
7. Bora, R.R., Lei, M., Tester, J.W., Lehmann, J., & You, F. (2020). Life cycle assessment and technoeconomic analysis of thermochemical conversion technologies applied to poultry litter with energy and nutrient recovery. *ACS Sustain. Chemistry Engineering* 8(22). <https://doi.org/10.1021/acssuschemeng.0c02860>.
8. Bordin, F.M., Linke, B.S., Fredel, M.C., Xavier, F.A., & Weingaertner, W.L. (2022). Grinding sustainability analysis using a fatigue failure-related product life indicator, *CIRP Journal of Manufacturing Science and Technology* 38: 1-15.
9. Bright, A. (2008). Vocalizations and acoustic parameters of flock noise from feather pecking and non-feather pecking laying flocks. *Br. Poultry Science* 49: 241249. <https://doi.org/10.1080/00071660802094172>.
10. Bumanis, N., Arhipova, I., Paura, L., Vitols, G., & Jankovska, L. (2022). *Data conceptual model for smart poultry farm management system*, 3rd International Conference on Industry 4.0 and Smart Manufacturing, 517-526.
11. Bustamante, E., Guijarro, E., García-Diego, F.J., Balasch, S., Hospitaler, A., & Torres, A.G. (2012). Multisensory system for is temporal measurements to assess indoor climatic conditions in poultry farms. *Sensors (Switzerland)* 12: 5752–5774. <https://doi.org/10.3390/s120505752>.
12. Calvet, S., Campelo, J., Estellés, F., Perles, A., Mercado, R., Serrano, J., Calvet, S., Campelo, J.C., Estellés, F., Perles, A., Mercado, R., & Serrano, J.J. (2014). Suitability evaluation of multipoint simultaneous CO₂ sampling wireless sensors for livestock buildings. *Sensors* 14: 10479–10496. <https://doi.org/10.3390/s140610479>.
13. Carroll, B.T., Anderson, D. V., Daley, W., Harbert, S., Britton, D.F., & Jackwood, M.W. (2014). *Detecting symptoms of diseases in poultry through audio signal processing*. 2014 IEEE Glob. Conf. Signal Inf. Process. 1132–1135. <https://doi.org/10.1109/GlobalSIP.2014.7032298>.

14. Chakchai, S., Mongkonchai, N., Aimtongkham, P., Wijitsopon, K., & Rujirakul, K. (2014). An evaluation of data-mining classification models for network intrusion detection. In 4th international conference on digital information and communication technology and it's applications (pp. 90–94). Bangkok.
15. Chen, J., Hu, T., Wu, J.H., Si, H.P., & Lin, K.Y. (2014). Applications of internet of things in facility agriculture. *Applied Mechanics and Materials* 685: 517–523. <https://doi.org/10.4028/www.scientific.net/AMM.685.517>.
16. Chen, L., & Neethirajan, S. (2015). A homogenous fluorescence quenching based assay for specific and sensitive detection of influenza virus *Ahemag glutininantigen*. *Sensors* (Basel) 15: 8852–8865. <https://doi.org/10.3390/s150408852>.
17. Chen, M., Mao, S., & Liu, Y. (2014). Big data: A survey. *Mob. Networks Appl.* 19: 171–209. <https://doi.org/10.1007/s11036-013-0489-0>.
18. Chowdhury, E.U., & Morey, A. (2019). Intelligent packaging for poultry industry, *Poultry Science Association Inc.* 791-800.
19. Colles, F.M., Cain, R.J., Nickson, T., Smith, A.L., Roberts, S.J., Maiden, M.C.J., Lunn, D., & Dawkins, M.S., (2016). Monitoring chicken flock behavior provides early warning of infection by human pathogen *Campylobacter*. *Proc. R. Soc. B Biology Science* 283: 1–6. <https://doi.org/10.1098/rspb.2015.2323>.
20. Comim, F., & Hirai, T. (2022). Sustainability and human development indicators: a poset analysis, *Ecological Economics* 198: 107470.
21. David, B., Mejdell, C., Michel, V., Lund, V., & Oppermann Moe, R. (2015). Air quality in alternative housing systems may have an impact on laying hen welfare. PartII—Ammonia. *Anim. Open Access Journal from MDPI* 5: 886–896. <https://doi.org/10.3390/ani5030389>.
22. De Longe, M.S., Miles, A., & Carlisle, L. (2016). Investing in the transition to sustainable agriculture. *Environmental Science & Policy* 55: 266-273.
23. Epp, M. (2019). Poultry Technology: Rise of the robots [WWW Document]. *Can. Poult.* URL <https://www.canadianpoultrymag.com/production/new-technology/rise-of-the-robots-30876?jij=1570740094606> (accessed 10.10.19).
24. FAO. (2017). Ending poverty and hunger by investing in agriculture and rural areas. Food and Agriculture Organization of the United Nations (FAO).
25. Fontana, I., Tullo, E., Butterworth, A., & Guarino, M. (2015). An innovative approach to predict the growth in intensive poultry farming. *Computer Electronic Agriculture* 119: 178–183. <https://doi.org/10.1016/j.compag.2015.10.001>.
26. GeSI. (2016). How Digital Solutions Will Drive Progress Towards The Sustainable Development Goals. Brussels: Global e-Sustainability Initiative (GeSI).
27. Ghasemi, R., Mohaghar, A., Safari, H. (2016). "Prioritization of IoT technology applications in the Iranian health sector: a stimulus for sustainable development", Faculty of Management, University of Tehran, 8(1). (In Persian)
28. Gunnarsdottir, J., Davidsdottir, B., Worrell, E., & Sigurgeirsdottir, S. (2022). Indicators for sustainable energy development: an Icelandic case study, *Energy Policy* 164: 112926.
29. Halachmi, I., Guarino, M., & Bewley, J. (2019), Smart animal agriculture: application of real-time sensors to improve animal well-being and production. *Annu. Rev. Animal Bioscience* 7: 403–25
30. Huang, L., & Liu, Ch. (2016) "The Application Mode in Urban Transportation Management Based on Internet of Things", Proceedings of the 2nd International Conference on Computer Science and Electronics Engineering (ICCSE 2013) B. Nonnecke, M. Bruch," IoT & Sustainability: Practice, Policy and Promise", Center for Information Technology Research in the Interest of Society& the Banatao Institute University of California.

31. Hadinia, S.H., Carneiro, P.R.O., Ouellette, C.A., & Zuidhof, M.J. (2018). Energy partitioning by broiler breeder pullets in skip-a-day and precision feeding systems. *Poultry Science*. <http://doi.org/10.3382/ps/pey283>.
32. Information Builders. (2011). Seven Steps to Effective Data Governance: A White Paper.
33. Jackman, P., Ward, S., Brennan, L., & Corkery, G. (2015). Application of wireless technologies to forward predict crop yields in the poultry production chain. *CIGR Journal* 17: 287–295.
34. Kamilaris, A., Kartakoullis, A., & Prenafeta-Boldú, F.X. (2017). A review on the practice of big data analysis in agriculture. *Computer Electronic Agriculture* 143: 23–37. <https://doi.org/10.1016/J.COMPAG.2017.09.037>.
35. Kesharavz-Ghorabae, M., Amiri, M., Zavadskas, E.K., Turshis, Z., & Antucheviciene, J. (2018). Simultaneous evaluation of criteria and alternatives (SECA) for multi-criteria decision-making. *INFORMATICA* 29(2): 265-280.
36. Kocovic, P., Behringer, R., Ramachandran, M., & Mihajlovic, R. (2017). Emerging Trends and Applications of the Internet of Things. Hershey, PA: IGI Global. <http://doi.org/10.4018/978-1-5225-2437-3>.
37. Manika, J., Chui, M., Brown, B., Bughin, J., Dobbs, R., & Byers, A.H. (2011). Big Data: The Next Frontier for Innovation, Competition, and Productivity. McKinsey Global Institute.
38. Marchewka, J., Watanabe, T.T.N., Ferrante, V., & Estevez, I. (2013). Welfareassessmentinbroiler farms: Transect walks versus individual scoring. *Pollution Science* 92: 2588–2599. <https://doi.org/10.3382/ps.2013-03229>.
39. Manteuffel, G., Puppe, B., & Schön, P.C. (2004). Vocalization of farm animals as a measure of welfare. *Appl. Animal Behavior Science* 88: 163–182. <https://doi.org/10.1016/j.applanim.2004.02.012>.
40. Merad, M., Dechy, N., Serir, L., Grabisch, M., & Marcel, F. (2013). Using a multi-criteria decision aid methodology to implement sustainable development principles within an organization. *European Journal of Operational Research* 224(3): 603-613. <http://doi.org/10.1016/j.ejor.2012.08.019>.
41. McDougal, T. (2018). Poultry World - French poultry robot up and running, [https://www.poultryworld.net/Meat/Articles/2018/1/Frenchpoultry-robotup-and-running-231202E/\(accessed 1.22.19\)](https://www.poultryworld.net/Meat/Articles/2018/1/Frenchpoultry-robotup-and-running-231202E/(accessed%201.22.19)).
42. Mohammadian, A., Heidari Dehvaei, J., & Ghorbani, A. (2020). Prioritize IoT applications in agriculture using sustainable development indicators. *Iranian Agricultural Economics and Development Research* 51(4): 759-745. (Persian)
43. Mohammadzadeh, A., Ghafoori, S., Mohammadian, A., Mohammadkazemi, R., Mahbanooei, B., & Ghasemi, R. (2018). A Fuzzy Analytic Network Process (FANP) approach for prioritizing internet of things challenges in Iran. *Technology in Society* 53: 124-134. <http://doi.org/10.1016/j.techsoc.2018.01.007>.
44. Nääs, I.A., Garcia, R.G., & Caldara, F.R. (2014). Infrared thermal image for assessing animal health and welfare. *Journal Animal Behavior Biometeorology* 2: 66–72. <https://doi.org/10.14269/2318-1265/jabb.v2n3p66-72>.
45. Neethirajan, S., Tuteja, S., Huang, S.-T., & Kelton, D. (2017). Recent advances in wearable sensors for animal and livestock health management. *Biosens. Bioelectron* 98: 398–407. <https://doi.org/10.1016/j.sbsr.2016.11.004>.
46. Nuñez, I.A., & Ross, T.M. (2019). A review of H5Nx avian influenza viruses. *Ther. Adv. Vaccin. Immunother.* 7: 2515135518821625. <http://doi.org/10.1177/2515135518821625>.
47. Luka, G., Ahmadi, A., Najjaran, H., Alocilja, E., Derosa, M., Wolthers, K., Malki, A., Aziz, H., Althani, A., & Hoorfar, M. (2015). Microfluidics integrated biosensors: A leading

- technology towards lab-on-A-chip and sensing applications. *Sensors (Switzerland)* 15: 30011–30031. <https://doi.org/10.3390/s151229783>.
48. Obedient Langroudi, S.H., & Shamsai, E. (2007). Rural development based on agricultural continuity and sustainability, *Geographical Research Quarterly* 86: 85-104. (In Persian)
 49. Okada, H., Itoh, T., Suzuki, K., & Tsukamoto, K. (2009). Wireless sensor system for detection of avian influenza outbreak farms at an early stage. *IEEE Sensors* 1374–1377. <https://doi.org/10.1109/ICSENS.2009.5398422>.
 50. Okada, H., Suzuki, K., Kenji, T., & Itoh, T. (2014). Applicability of wireless activity sensor network to avian influenza monitoring system in poultry farms. *Journal Sens. Technology* 4: 18–23.
 51. O'Grady, M.J., & O'Hare, G.M.J.I. (2017). Modelling the smart farm. 4(3): 179-187.
 52. Park, M., Britton, D., Daley, W., MsMurray, G., Navaei, M., Samoylov, A., Usher, C., & Xu, J. (2022). Artificial intelligence, sensors, robots, and transportation systems drive an innovative future for poultry broiler and breeder management, *Animal Frontiers* 12(2): 40-48.
 53. Poursaeed, A., Mirdamadi, M., Malekmohammadi, I., & Hosseini, J. (2010). The partnership models of agricultural sustainable development based on Multiple Criteria Decision Making (MCDM) in Iran. *African Journal of Agricultural Research* 5(23): 3185-3190.
 54. Praveen, K., & Satish, C. (2012). An intelligent system based on sensor integration & sensor fusion. *International Journal of Advanced Research in Computer Science and Software Engineering* 2(3): 2277–2283.
 55. Quaddus, M., & Siddique, M. (2001). Modelling sustainable development planning: A multicriteria decision conferencing approach. *Environment International* 27(2-3): 89-95. [http://doi.org/10.1016/S0160-4120\(01\)00066](http://doi.org/10.1016/S0160-4120(01)00066).
 56. Quintero-Angel, M., & González-Acevedo, A. (2018): Tendencies and challenges for the assessment of agricultural sustainability. *Agriculture, Ecosystems & Environment* 254: 273-281.
 57. Regan, Á. (2019). 'Smart farming' in Ireland: A risk perception study with key governance actors. *NJAS-Wageningen Journal of Life Sciences* 90: 100292.
 58. Rezaei-Moghaddam, K., & Karami, E. (2008). A multiple criteria evaluation of sustainable agricultural development models using AHP. *Sustainable Agricultural Development Models* 10(4): 407-426.
 59. Rizwan, M., Carroll, B.T., Anderson, D.V., Daley, W., Harbert, S., Britton, D.F., & Jackwood, M.W. (2017). *Identifying rale sounds in chickens using audio signals for early disease detection in poultry*. 2016 IEEE Glob. Conf. Signal Inf. Process. Glob. 2016 - Proc. 55–59. <https://doi.org/10.1109/GlobalSIP.2016.7905802>.
 60. Ronald, B., & Siarhei, S. (2012). Experiments for evaluating sensors' precision in wireless sensor networks. *International Journal of Ad Hoc, Sensor & Ubiquitous Computing* 3(3): 105–121.
 61. Rani, K.S.S., & Devarajan, N. (2012). Multi objective sensor node deployment in wireless sensor networks. *International Journal of Engineering Science and Technology* 4(4): 1262–1266.
 62. Sandelowski, M., & Barroso, J. (2006). *Handbook for Synthesizing Qualitative Research*. New York: Springer Publishing Company, Inc.
 63. Sassi, N. Ben, Averós, X., Estevez, I. (2016). Technology and poultry welfare. *Animals* 6: 1–21. <https://doi.org/10.3390/ani6100062>.
 64. Schuetz, C.G., Schausberger, S., & Schrefl, M. (2018). Building an active semantic data ware house for precision dairy farming. *Journal Organ. Computer Electronic Commer.* 28: 122–141. <https://doi.org/10.1080/10919392.2018.1444344>.

65. Senoret, A., Ramirez, M., & Rehner, J. (2022). Employment and sustainability: the relation between precarious work and spatial inequality in the neoliberal city, *World Development* 153: 105840.
66. Shang, X. (2012). "Internet of Things (IOT) Service: Architecture and its application in E-Commerce", *Journal of Electronic Commerce in Organizations* 10(3): 44-55.
67. Shinder, D., Rusal, M., Giloh, M., & Yahav, S. (2009). Effect of repetitive acute cold exposures during the last phase of broiler embryogenesis on cold resistance through the lifespan. *Poult. Science* 88: 636–646. <https://doi.org/10.3382/ps.2008-00213>.
68. Sicular, S. (2013). Gartner's Big Data Definition Consists of Three Parts, Not to Be Confused with Three "V"s [WWW Document]. URL <https://www.forbes.com/sites/gartnergroup/2013/03/27/gartners-big-datadefinition-consists-of-three-parts-notto-be-confused-with-threevs/#4912508b42f6> (accessed 5.22.18).
69. Silvera, A.M., Knowles, T.G., Butterworth, A., Berckmans, D., Vranken, E., & Blokhuis, H.J. (2017). Lameness assessment with automatic monitoring of activity in commercial broiler flocks. *Poult. Science* 96: 2013–2017. <https://doi.org/10.3382/ps/pex023>.
70. Smith, D., Lyle, S., Berry, A., Manning, N., Zaki, M., & Neely, A. (2015). Internet of animal health things opportunities and challenges data and analytics: Internet of Animal Health Things. <https://doi.org/10.13140/RG.2.1.1113.8409>.
71. Sun, Chunling. (2012). "Application of RFID Technology for Logistics on Internet of Things ", AASRI Conference on Computational Intelligence and Bioinformatics, Published by Elsevier.
72. Vanderhasselt, R.F., Sprenger, M., Duchateau, L., & Tuytens, F.M. (2013). Automated assessment of footpad dermatitis in broiler chickens at the slaughter-line: Evaluation and correspondence with human expert scores. *Poult. Science* 92: 12–8. <https://doi.org/10.3382/ps.2012-02153>.
73. Veisi, H., Liaghati, H., & Alipour, A. (2016). Developing an ethics-based approach to indicators of sustainable agriculture using analytic hierarchy process (AHP). *Ecological Indicators* 60: 644-654.
74. Verma, N., & Usman, A. (2016). Internet of Things (IoT): A relief for Indian farmers. 2016 IEEE Global Humanitarian Technology Conference (GHTC) (pp. 831-835). Seattle: IEEE.
75. Walters, S.J. (2009). *Quality of Life Outcomes in Clinical Trials and Health-Care Evaluation: A Practical Guide to Analysis and Interpretation*. Wiley, New York.
76. Walter, A., Finger, R., Huber, R., & Buchmann, N. (2017). Opinion: Smart farming is key to developing sustainable agriculture. *Proceedings of the National Academy of Sciences* 114(24): 6148–6150. <http://doi.org/10.1073/pnas.1707462114>.
77. Wang, T.-C., & Lee, H.-D. (2009). Developing a fuzzy TOPSIS approach based on subjective weights and objective weights. *Expert Systems with Applications* 36(5): 8980–8985.
78. Wang, Y., & Qi, H. (2012). "Research of intelligent transportation system based on the internet of things frame", *Wireless Engineering and Technology* 3(3): 21471. <http://doi.org/10.4236/wet.2012.33023>.
79. Weersink, A., Fraser, E., Pannell, D., Duncan, E., & Rotz, S. (2018). Opportunities and challenges for big data in agricultural and environmental analysis. *Annual Review of Resource Economics* 10: 19-37.
80. Williams, J., Alter, T., & Shrivastava, P. (2018). Systemic governance of sustainable agriculture: Implementing sustainable development goals and climate-friendly farming. *Outlook on Agriculture* 47(3): 192-195. <http://doi.org/10.1177/0030727018795907>.
81. Xin, J., & Zazueta, F.J.A.E.I.C.J. (2016). Technology trends in ICT—towards data-driven, farmer centered and knowledge-based hybrid cloud architectures for smart farming 18(4): 275-279.

82. Xin, J., & Zazueta, F. (2016). Technology trends in ICT—towards data-driven, farmer-centered and knowledge-based hybrid cloud architectures for smart farming. *Agricultural Engineering International: CIGR Journal* 18(4): 275-279.
83. Xin, H., & Liu, K. (2017). Precision livestock farming in egg production. *Animal Front.* 7: 24. <https://doi.org/10.2527/af.2017.0105>.
84. Wolfert, S., Ge, L., Verdouw, C., & Bogaardt, M.J. (2017). Big data in smart farming—a review. *Agricultural Systems* 153: 69-80.
85. Zarei, M., Mohammadian, A., & Ghasemi, R. (2016). Internet of things in industries: a survey for sustainable development. *Innovation and Sustainable Development* 10(4): 419-442.
86. Zhang, Q., Yang, X.L., Zhou, Y.M., Wang, L.R., Guo, X.S. (2007). A wireless solution for greenhouse monitoring and control system based on Zig Bee technology. *Journal of Zhejiang University Science* 8(10): 1584–1587.
87. Zuidhof, M.J. (2018). Lifetime productivity of conventionally and precision-fed broiler breeders. *Poult. Science* In Press. <https://doi.org/10.3382/ps>.
88. Zuidhof, M.J., Fedorak, M.V., Ouellette, C.A., & Wenger, I.I. (2017). Precision feeding: Innovative management of broiler breeder feed intake and flock uniformity. *Poult. Science* 96: 2254–2263. <https://doi.org/10.3382/ps/pex013>.



مقاله پژوهشی

جلد ۳۶ شماره ۴، زمستان ۱۴۰۱، ص. ۴۱۱-۳۹۳

شناسایی و اولویت‌بندی راه‌حل‌های هوشمند در صنعت طیور براساس معیارهای پایداری

آمنه خدیور^{۱*} - فاطمه مجیبیان^۲ - زهرا ترکاشوند^۳

تاریخ دریافت: ۱۴۰۱/۰۳/۲۱

تاریخ پذیرش: ۱۴۰۱/۰۶/۲۳

چکیده

یکی از بخش‌های غذایی که تولید هر چه بیشتر در آن می‌تواند ضمن از بین بردن وابستگی به خارج صادرات ارزآوری نیز داشته باشد، بخش تولید و عرضه دام و طیور است. مرغداری صنعت مهمی برای تأمین پایدار غذا در کشورها می‌باشد. در این پژوهش کاربردها و راه‌حل‌های هوشمند در صنعت طیور شناسایی شده و با استفاده از شاخص‌های توسعه پایدار و با بهره‌گیری از روش ارزیابی همزمان معیارها و گزینه‌ها (SECA) به اولویت‌بندی این کاربردها پرداخته شده است. بر اساس تحلیل‌های صورت گرفته ۱۸ حوزه اصلی از راه‌حل‌های هوشمند در صنعت مرغداری شناسایی گردیده است. اوزان شاخص‌های توسعه پایدار بر اساس روش SECA، اقتصادی (۰/۳۵۱)، اجتماعی (۰/۳۳۸۳) و زیست محیطی (۰/۳۰۶۵) نشان می‌دهد که برای پیاده سازی پروژه‌های مبتنی بر راه‌حل‌های هوشمند در صنعت مرغداری بایستی بیشتر به پایداری اقتصادی اهمیت داد. این در حالیست که یکی از چالش‌های اصلی که بخش کشاورزی بخصوص صنعت مرغداری با آن روبرو است استفاده از روش‌های تولید کشاورزی سنتی است که باعث شده بیش از ظرفیت زمین استفاده کند. علاوه بر آن جهانی شدن، تغییرات آب و هوایی، حرکت از اقتصاد مبتنی بر سوخت فسیلی به سوی اقتصاد مبتنی بر محیط‌زیست و رقابت بر سر زمین، آب شیرین و نیروی کار منجر به پیچیدگی‌های بیشتر و ایجاد چالش تغذیه در جهان و بهره‌برداری اضافی شده است. با توجه به پتانسیل بالقوه راه‌حل‌های هوشمند در تحقق اهداف توسعه پایدار، پیشنهاد می‌شود که بیشتر بر روی جنبه زیست‌محیطی پروژه‌ها تمرکز شود.

واژه‌های کلیدی: اینترنت اشیا، توسعه پایدار، راه‌حل‌های هوشمند، روش SECA، صنعت طیور

۱ و ۳- به ترتیب دانشیار گروه مدیریت و کارشناس ارشد، دانشگاه الزهراء، تهران، ایران

*- نویسنده مسئول: (Email: a_khadivar@yahoo.com)

۲- دکترای تخصصی، دانشکده مدیریت و اقتصاد، دانشگاه تربیت مدرس